

GUIDE TO FULFILLING VIVITROL **PRESCRIPTIONS** **AND PATIENT** **SUPPORT SERVICES**

Please see [Important Safety Information](#) on pages 6 and 7.
Please see [Prescribing Information](#) and [Medication Guide](#).
Review [Medication Guide](#) with your patients.

Vivitrol[®]
(naltrexone for extended-release
injectable suspension)

ORDERING

VIVITROL® (naltrexone for extended-release injectable suspension)

You have the option to use Vivitrol2getherSM dedicated case manager services

Vivitrol2gether Patient Support Services is designed to help with:

- Reimbursement coverage verification
- Submission of VIVITROL prescriptions to a pharmacy (specialty or other)
- Submission of VIVITROL® Co-pay Savings Program application for eligible patients*
- Providing patient support services throughout fulfillment, transition of care, and during treatment with VIVITROL, through a dedicated Nurse Coordinator



For assistance with fulfilling VIVITROL prescriptions, fax ALL of the following documents to Vivitrol2gether Patient Support Services at 1-877-329-8484:

- Vivitrol2gether Enrollment Form (download an editable PDF at VIVITROLHCP.com/support)
- Photocopy of front/back of patient's insurance card (enlarged for legibility)
- Prior Authorization (PA) form (if required)

Sample Vivitrol2gether Enrollment Form

Our team is ready to support you and your patients.
Call 1-800-VIVITROL (1-800-848-4876) 9am–8pm (EST) and visit VIVITROLHCP.com/support to access helpful resources.

CRITICAL!

- Fax the PA form (if required) and an enlarged photocopy of patient's pharmacy and health insurance cards to the health insurer's toll-free fax number on the PA form. Pharmacies and health insurers typically prefer that these come directly from your office
- Follow up within 24 hours to ensure the prescription is being processed. Remember to identify date of scheduled injection. If the pharmacy cannot process the VIVITROL prescription, consider enrolling your patient into Vivitrol2gether to receive assistance with pharmacy routing

*Eligibility for Alkermes-Sponsored Co-pay Assistance: Offer valid only for prescriptions for FDA-approved indications. You must be at least 18 years old. If you are purchasing your VIVITROL prescriptions with benefits from Medicare, including Medicare Part D or Medicare Advantage plans; Medicaid, including Medicaid Managed Care or Alternative Benefit Plans ("ABPs") under the Affordable Care Act; Medigap; Veterans Administration ("VA"); Department of Defense ("DoD"); TriCare®; or any similar state funded programs such as medical or pharmaceutical assistance programs, you are not eligible for this offer. Void where prohibited by law, taxed, or restricted. Subject to plan benefit design requirements. Alkermes, Inc. reserves the right to rescind, revoke, or amend these offers without notice.

Alkermes® Contact(s)

Please see **Important Safety Information** on pages 6 and 7.
Please see **Prescribing Information** and **Medication Guide**.
Review **Medication Guide** with your patients.

Work directly with a specialty pharmacy (SP)

1 FIRST VIVITROL PRESCRIPTION	2 SP VERIFICATION
<p>TO DO</p> <ul style="list-style-type: none"> ▪ Determine available SP options with your representative ▪ Fax identified medical and pharmacy insurance cards and enrollment form to the SP ▪ Give the patient a copy of the enrollment form and keep one for their chart ▪ Work with the SP to complete the PA if the plan requires one 	<ul style="list-style-type: none"> ▪ SP verifies benefits; confirms prescription* ▪ Confirm status of PA if submitted, or start PA process with provider if not yet submitted ▪ SP applies co-pay savings to eligible patients
<p>CHECKLIST</p> <ul style="list-style-type: none"> ✓ Photocopies of the front and back of the medical and pharmacy insurance cards are enlarged to ensure legibility ✓ Enrollment form is legible ✓ All patient or provider signatures are provided ✓ Injection provider section is complete ✓ Contact and alternate contact information is provided 	<ul style="list-style-type: none"> ✓ To help avoid delays, provider must return SP calls if additional information is required ✓ If eligible, be sure co-pay savings is applied
<p>TIPS</p> <ul style="list-style-type: none"> ▪ Be sure all forms are complete and legible. Enlarge photocopies to ensure readability ▪ Be sure health plan and SP have a copy of completed PA form (if required) 	<ul style="list-style-type: none"> ▪ Follow up regularly with SP via phone, email, or fax ▪ Provide contact information to the SP, including primary/secondary phone and email addresses ▪ The patient should have a copy of their co-pay card so they can remind the SP they are eligible for the co-pay program ▪ Remind the patient that the SP will contact them

Specialty Pharmacy Contact(s)

Please see **Important Safety Information** on pages 6 and 7. Please see **Prescribing Information** and **Medication Guide**. Review **Medication Guide** with your patients.



Work directly with a specialty pharmacy (SP)

3

SHIPMENT OF VIVITROL

4

DELIVERY AND
INJECTION OF VIVITROL

TO DO

- SP must contact patient (or financially responsible designee) to authorize billing and shipment
- Instruct patient to take or return SP call in order to authorize shipment
- Provide patient with SP name and phone numbers
- Once authorized, SP contacts the office to schedule shipment
- Confirm receipt of VIVITROL
- Patient receives injection of VIVITROL
- Schedule next injection
- Notify SP of next VIVITROL injection date
- Make sure the patient knows the next injection date and the SP name/phone number to authorize shipment

CHECKLIST

- ✓ Patient has authorized shipment or been provided with the SP number to call
- ✓ Alternate patient contacts provided
- ✓ Shipment is coordinated with the timing of the injection
- ✓ Create a discharge plan for the patient, including follow-on injection and care. For assistance, call 1-800-VIVITROL or consult VIVITROLHCP.com/support to locate an injection provider
- ✓ Communicate to patient that the SP will contact them every month to authorize shipment

TIPS

- Educate designees, if used, on VIVITROL
- Communicate required VIVITROL injection date for each patient
- Follow up with the patient to ensure the documentation is signed and returned
- Confirm patient contact information and notify SP of any changes

*If SP is not contracted to fill prescriptions for a specific insurer, the prescription is transferred to another pharmacy.


Please see **Important Safety Information** on pages 6 and 7.
Please see **Prescribing Information** and **Medication Guide**.
Review **Medication Guide** with your patients.

Vivitrol[®]
(naltrexone for extended-release
injectable suspension)

Purchase directly buy and bill programs

- 1 Contact payer to confirm HCPCS code J2315 is covered by patient's policy**
- 2 Register with Opus Health to become a co-pay card participant. Call 1-877-838-3836 for assistance. You'll receive a practice-populated form for enrollment**
After receiving the EOB, send the form and co-pay card to OPUS Health
- 3 Ask a VIVITROL representative for information on how to open an account with an Alkermes-preferred distribution partner**
- 4 Order VIVITROL and inject the patient**
- 5 Fill out and send the appropriate form below:**


Sample CMS-1500 Claim Form

A sample CMS-1500 Health Insurance Claim Form, showing various fields for patient information, provider details, and charges. A large 'SAMPLE' watermark is overlaid on the form.

Specify
380 mg


Enter HCPCS
code **J2315**
Enter CPT[†]
code **96372**

Sample CMS-1450 (UB-04) Claim Form

A sample CMS-1450 (UB-04) Institutional Claim Form, showing various fields for patient information, provider details, and charges. A large 'SAMPLE' watermark is overlaid on the form.

These documents are provided for your guidance only. Please contact the payer or Vivitrol2getherSM to verify coding and claim information.

For more information, download the Billing & Coding Resource at VIVITROLHCP.com/support



*Each company retains sole discretion for opening an account and establishing credit limits.
†CPT[®]=Current Procedural Terminology. Copyright of the American Medical Association, 2018.
EOB=explanation of benefits; HCPCS=Healthcare Common Procedure Coding System.

Please see **Important Safety Information** on pages 6 and 7.
Please see **Prescribing Information** and **Medication Guide**.
Review **Medication Guide** with your patients.

Vivitrol[®]
(naltrexone for extended-release
injectable suspension)

IMPORTANT SAFETY INFORMATION

FOR VIVITROL[®] (naltrexone for extended-release injectable suspension)

INDICATIONS

VIVITROL is indicated for:

- Treatment of alcohol dependence in patients who are able to abstain from alcohol in an outpatient setting prior to initiation of treatment with VIVITROL. Patients should not be actively drinking at the time of initial VIVITROL administration.
- Prevention of relapse to opioid dependence, following opioid detoxification.
- VIVITROL should be part of a comprehensive management program that includes psychosocial support.

CONTRAINDICATIONS

VIVITROL is contraindicated in patients:

- Receiving opioid analgesics
- With current physiologic opioid dependence
- In acute opioid withdrawal
- Who have failed the naloxone challenge test or have a positive urine screen for opioids
- Who have exhibited hypersensitivity to naltrexone, polylactide-co-glycolide (PLG), carboxymethylcellulose, or any other components of the diluent

WARNINGS/PRECAUTIONS

Vulnerability to Opioid Overdose:

- After opioid detoxification, patients are likely to have a reduced tolerance to opioids. VIVITROL blocks the effects of exogenous opioids for approximately 28 days after administration. As the blockade wanes and eventually dissipates completely, use of previously tolerated doses of opioids could result in potentially life-threatening opioid intoxication (respiratory compromise or arrest, circulatory collapse, etc.).
- Cases of opioid overdose with fatal outcomes have been reported in patients who used opioids at the end of a dosing interval, after missing a scheduled dose, or after discontinuing treatment. Patients and caregivers should be told of this increased sensitivity to opioids and the risk of overdose.
- Although VIVITROL is a potent antagonist with a prolonged pharmacological effect, the blockade produced by VIVITROL is surmountable. The plasma concentration of exogenous opioids attained immediately following their acute administration may be sufficient to overcome the competitive receptor blockade. This poses a potential risk to individuals who

attempt, on their own, to overcome the blockade by administering large amounts of exogenous opioids.

- Any attempt by a patient to overcome the VIVITROL blockade by taking opioids may lead to fatal overdose. Patients should be told of the serious consequences of trying to overcome the opioid blockade.

Injection Site Reactions:

- VIVITROL injections may be followed by pain, tenderness, induration, swelling, erythema, bruising, or pruritus; however, in some cases injection site reactions may be very severe.
- Injection site reactions not improving may require prompt medical attention, including, in some cases, surgical intervention.
- Inadvertent subcutaneous/adipose layer injection of VIVITROL may increase the likelihood of severe injection site reactions.
- Select proper needle size for patient body habitus, and use only the needles provided in the carton.
- Patients should be informed that any concerning injection site reactions should be brought to the attention of their healthcare provider.

Precipitation of Opioid Withdrawal:

- When withdrawal is precipitated abruptly by administration of an opioid antagonist to an opioid-dependent patient, the resulting withdrawal syndrome can be severe. Some cases of withdrawal symptoms have been severe enough to require hospitalization, and in some cases, management in the ICU.
- To prevent occurrence of precipitated withdrawal, opioid-dependent patients, including those being treated for alcohol dependence, should be opioid-free (including tramadol) before starting VIVITROL treatment:
 - An opioid-free interval of a minimum of 7-10 days is recommended for patients previously dependent on short-acting opioids.
 - Patients transitioning from buprenorphine or methadone may be vulnerable to precipitated withdrawal for as long as two weeks.
- If a more rapid transition from agonist to antagonist therapy is deemed necessary and appropriate by the healthcare provider, monitor the patient closely in an appropriate medical setting where precipitated withdrawal can be managed.

IMPORTANT SAFETY INFORMATION (cont'd)

FOR VIVITROL[®] (naltrexone for extended-release injectable suspension)

- Patients should be made aware of the risk associated with precipitated withdrawal and be encouraged to give an accurate account of last opioid use.

Hepatotoxicity:

- Cases of hepatitis and clinically significant liver dysfunction have been observed in association with VIVITROL. Warn patients of the risk of hepatic injury; advise them to seek help if experiencing symptoms of acute hepatitis. Discontinue use of VIVITROL in patients who exhibit acute hepatitis symptoms.

Depression and Suicidality:

- Alcohol- and opioid-dependent patients taking VIVITROL should be monitored for depression or suicidal thoughts. Alert families and caregivers to monitor and report the emergence of symptoms of depression or suicidality.

When Reversal of VIVITROL Blockade Is Required for Pain Management:

- For VIVITROL patients in emergency situations, suggestions for pain management include regional analgesia or use of non-opioid analgesics. If opioid therapy is required to reverse the VIVITROL blockade, patients should be closely monitored by trained personnel in a setting staffed and equipped for CPR.

Eosinophilic Pneumonia:

- Cases of eosinophilic pneumonia requiring hospitalization have been reported. Warn patients of the risk of eosinophilic pneumonia and to seek medical attention if they develop symptoms of pneumonia.

Hypersensitivity Reactions:

- Patients should be warned of the risk of hypersensitivity reactions, including anaphylaxis.

Intramuscular Injections:

- As with any intramuscular injection, VIVITROL should be administered with caution to patients with thrombocytopenia or any coagulation disorder.

Alcohol Withdrawal:

- Use of VIVITROL does not eliminate nor diminish alcohol withdrawal symptoms.

ADVERSE REACTIONS

- The adverse events seen most frequently in association with VIVITROL therapy for alcohol dependence (ie, those occurring in $\geq 5\%$ and at least twice as frequently with VIVITROL than placebo) include nausea, vomiting, injection site reactions (including induration, pruritus, nodules, and swelling), arthralgia, arthritis, or joint stiffness, muscle cramps, dizziness or syncope, somnolence or sedation, anorexia, decreased appetite or other appetite disorders.
- The adverse events seen most frequently in association with VIVITROL in opioid-dependent patients (ie, those occurring in $\geq 2\%$ and at least twice as frequently with VIVITROL than placebo) were hepatic enzyme abnormalities, injection site pain, nasopharyngitis, insomnia, and toothache.

You are encouraged to report side effects to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Prescribing Information and Medication Guide. Review Medication Guide with your patients.

KEY CONSIDERATIONS WHEN ORDERING VIVITROL[®] (naltrexone for extended-release injectable suspension)

Most important: ensure patients (and alternate contacts) understand the use of VIVITROL during their treatment and are prepared to return and receive calls as needed.

Make sure that:

- If VIVITROL is determined to be appropriate treatment, start planning as soon as possible
- All forms and photocopies of front/back of insurance cards are completely legible
- All patient and provider signatures are provided
- The health plan and SP have copies of the completed PA form, if required
- The SP has all relevant contact information
- The patient's contact/alternate contact information is satisfactory
- The patient has been given the SP contact information and instructed to take or return the SP call in order to authorize shipment of VIVITROL
- The next VIVITROL injection date is scheduled with the patient and SP and is communicated to the SP
- For buy and bill, the payer-confirmed code J2315 is covered by patient's policy and CPT[®] code 96372 and 380 mg are entered correctly on the form

*CPT[®]=Current Procedural Terminology.

Suggestions to help improve fill time

- Learn which pharmacies work with your patients' health plans
- Follow up regularly with SP contact person via phone, email, or fax
- Communicate required VIVITROL injection date for each patient



Please see **Important Safety Information** on pages 6 and 7.
Please see **Prescribing Information** and **Medication Guide**.
Review **Medication Guide** with your patients.



ALKERMES and VIVITROL are registered trademarks of Alkermes, Inc.
©2019 Alkermes, Inc. All rights reserved.
VIV-004207-v2
vivitrol.com

Vivitrol[®]
(naltrexone for extended-release injectable suspension)